

Post-rift seaward downwarping at passive margins: New insights from southern Oman using stratigraphy to constrain apatite fission-track and (U-Th)/He dating

Yanni Gunnell

Laboratoire de Géographie Physique, Université Paris 7, 2 place Jussieu, 75251 Paris Cedex 5, France

Andy Carter

School of Earth Science, Birkbeck College, University of London, Malet Street, London WC1E 7HX, UK

Carole Petit

Marc Fournier

Laboratoire de Tectonique, Université Paris 6, 4 place Jussieu, 75252 Paris Cedex 5, France

ABSTRACT

The plateau edge of southern Oman is used as a natural laboratory to independently test apatite fission-track analysis, (U-Th)/He dating, and stratigraphy as methods for quantifying post-rift erosion depths and lithospheric response at passive margins. A mappable unconformity between the Proterozoic basement and the low-conductivity, pre-rift sediment cover links residual buttes preserved at the coast to the escarpment, and therefore imposes tight limitations on parameter choices for modeling the Cenozoic topographic evolution. A mean geothermal gradient of $\sim 33^\circ\text{C km}^{-1}$ ensures the best fit between the thermochronologic data and the stratigraphically defined coast-to-scarp maximum denudation depth of ~ 1.75 km. With a lithospheric elastic thickness of 7 km, the resulting flexural response generates a retreat and uplift of the escarpment associated with a seaward downwarp of the unconformity, made possible because denudation minima are inboard of the escarpment and sediment loading occurs offshore. This closed experiment confirms the value of low-temperature thermochronology as a tool for quantifying long-term erosion, but also highlights the benefits of including independent stratigraphic information in terms of obtaining more plausible solutions. This study suggests that post-rift downwarping of eroded rift shoulders can occur, and may have not been commonly detected because low-temperature thermochronology data sets lack both suitable geologic data and intrinsic sensitivity.

Keywords: thermochronology, rifted margin, denudation, flexure, Oman.

INTRODUCTION

Low-temperature thermochronology (LTT) is commonly used to detect the depth and timing of erosion on 10–100 m.y. time scales at rifted continental margins (e.g., Braun and van der Beek, 2004), but cooling histories are indexed on a thermal reference frame within the Earth's crust. Conversion to a stratigraphic and topographic frame is therefore underpinned by an array of assumptions. In most models of rifted margin evolution, (1) the geometry of initial topography is arbitrary because information on initial and boundary conditions is lacking; (2) the value of pre-rift continental freeboard is speculative; and (3) estimates of lithospheric elastic thickness (EET) are indexed on the response to erosional unloading of the poorly defined initial topographic parameters. Sensitivity to initial conditions is often tested by letting geometries and EET vary, but solutions will be nonunique.

Depths of erosion control the isostatic response of the plateau edge to unloading, and are routinely obtained by thermal modeling of apatite fission track (AFT) data, more recently

in combination with apatite (U-Th)/He (hereafter AHe) ages. Assumptions made while converting rock cooling rates to denudation rates typically deal with nonunique rock-cooling paths and uncertainties in geothermal gradients. This further affects estimates of the magnitude and timing of denudation, critical in determining appropriate isostatic response functions of the lithosphere to unloading. In summary, long-term passive margin evolution models are sensitive to parameter choices, which profoundly affect our understanding of how landscapes evolve (Gunnell et al., 2003).

Optimal solutions can nevertheless be significantly narrowed when limitations afforded by independent data exist. This study focuses on southern Oman, where robust stratigraphic data suited to reconstructing long-term topographic evolution are available. Given the parameter sensitivity of LTT modeling, the objective here was to assess how well LTT can independently (1) provide matching estimates of the stratigraphically obtained denudation depths, and (2) reveal the coast-to-scarp pattern of denudation.

METHOD: STRATIGRAPHIC APPROACH

The southern edge of the Arabian plateau is a nonvolcanic, obliquely rifted margin (Fournier et al., 2004; Bellahsen et al., 2006). The plateau terminates as the south-facing Jabal Samhan escarpment (Fig. 1), rising to elevations of 1.8 km. It consists of an ~ 1 -km-thick sediment package tilted northward by a few degrees and overlying the Proterozoic basement. The age of rifting is not precisely established (see the GSA Data Repository¹), but extensional tectonics occurred after 35 Ma (Watchorn et al., 1998) and before the onset of seafloor spreading in the eastern Gulf of Aden at 18 Ma (Leroy et al., 2004). Continental freeboard afforded by initial shoulder

¹GSA Data Repository item 2007160, details on flexural and thermal modeling, low-temperature thermochronology analytic procedures, and related figures and data tables, is available online at www.geosociety.org/pubs/ft2007.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.




Figure 1. Southern Dhofar. A: Study area. **B:** Jabal Samhan (source: Google Earth). **C:** Topography, bathymetry (from Leroy et al., 2004), and main faults (master fault, MF, after D'Acremont et al., 2005). **D:** Lithology (after Mercolli et al., 2006, simplified) and apatite sample location (S1–S9). White arrows show buttes.

uplift provided the energy for antidip streams to drive scarp evolution (Petit et al., 2007).

Key stratigraphic details on the denudation history of Jabal Samhan are summarized here. (1) The conformable Eocene marine sequence capping the plateau allows calculation of vertical uplift, eroded thicknesses, and fault throws. The pre-rift paleotopography coincided with the top of the youngest Eocene marine deposit, which remains uneroded ~100 km inland from the coastline, where it was uplifted to elevations of ~0.6 km above sea level and preserved (Fig. 2). (2) The African-Arabian land surface at the Cretaceous-Tertiary boundary was a low-lying erosional plain capped by thin sedimentary Paleozoic and Cretaceous rocks, subsequently inundated and buried below sea level

by the Cenozoic marine sequence. The corresponding stratigraphic unconformity is continuously exposed at the escarpment. It can be used to track crustal deformation caused by rifting, erosion, isostasy, basement burial depths, and potential thermochronometric signatures. The coastal plain is entirely cut in basement rocks, which have been eroded to some depth below the basal unconformity due to rift-flank uplift. Buttes near the coastline retain the sedimentary caprock of the escarpment (Figs. 1 and 2); their presence is critical to this study. Detailed mapping (Platel et al., 1992; Mercolli et al., 2006) indicates that their preservation is not afforded by localized downfaulting, but rather they have survived because of limited erosion at those sites. Because the unconformity can be




Figure 2. Strike-perpendicular cross section of Jabal Samhan based on Platel et al. (1992). MF—master fault.

projected seaward from the escarpment to the buttes (Fig. 2), these provide important geometric data points for reconstructing erosion depths. A numerical model of the Dhofar margin (Petit et al., 2007), in which peak denudation depths in the inner coastal belt and sediment loading offshore explain the observed seaward flexural deformation of the margin, has provided EET values of 7 ± 1 km based on simultaneous best fits to the topography and the unconformity.

METHOD: RADIOMETRIC APPROACH

The nonradiometric reconstruction stands alone (Petit et al., 2007), but serves here as a reference frame against which to assess the accuracy of radiometrically derived denudation rates routinely used in passive margin settings (Gallagher et al., 1998). Nine samples (Fig. 1D) were collected across the coastal plain for LTT analysis in order to quantify denudation (analytic procedure in the Data Repository). AFT and AHe partial annealing and retention zones (PAZ and PRZ) span a range of temperatures with partially overlapping sensitivity windows. For low chlorine apatites, this approach can reconstruct thermal histories between ~120 °C and ~40 °C. AFT and AHe ages can be interpreted as cooling ages if cooling is monotonic and fairly rapid, because annealing and diffusive losses remain limited. When cooling is slower or locally complicated, only apparent AFT and AHe ages can be defined. To understand the significance of such data, modeling packages (e.g., Ketchum, 2005) are used to define probable thermal histories. Because thermal modeling is often used in settings where independent constraints on paleoburial depths are unknown, LTT studies predominantly use random searches of optimal constraints on time-temperature (t - T) histories permitted by the measured AFT and AHe data. The well-defined stratigraphic setting in Dhofar provides a rare opportunity for implementing forward modeling of the t - T data, a procedure in which thermal histories are fitted empirically to predefined independent geologic parameters. This approach predicts what combination of measured AFT and AHe parameters would be expected from a sample that has a predefined t - T history. It also affords indirect limitations on geothermal gradients in regions lacking heat flow data.

RESULTS

All AFT samples (Table DR1; see footnote 1) exhibit mean track lengths <14 μ m and a wide scatter in central ages (Fig. 3). None of the ages can therefore be interpreted as straightforward rapid cooling ages; this is typical of most passive margins. AFT ages are much older than the rifting age, which in combination with partially annealed track distributions (Fig. 3) suggests maximum paleoburial temperatures of 70–100 °C maintained over a relatively long period




Figure 3. Integrated low-temperature thermochronology results for southern DhoFar. Best-fitting cooling histories (top left) with track-length distributions (histogram) and their model fits (black curves). Cooling paths in gray used apatite fission track (AFT) and apatite (U-Th)/He (AHe) data simultaneously. For S6, AFT and AHe ages are close, and suggest comparatively rapid cooling through partial annealing zone. Undetected Cenozoic reheating event seems linked to sample's position on long-eroding southern footwall upland of Hasik graben (Fig. 1; Fournier et al., 2004).

of time. As such, the AFT signatures are typical of an outcrop of a PAZ. This is confirmed by the survival of short tracks in all the distributions. An extreme case is S9, where the bimodal track distribution and relatively young age suggest a complex thermal history.

Although only three samples (S4, S6, S9) yielded apatite crystals of suitable quality for AHe, these samples span the entire topographically uniform coastal plain. Discordant (U-Th)/He single-grain ages were handled by focusing on the mean age of the replicates (Fig. 3; Table DR2). These ages all point to rapid cool-

ing between 30 and 20 Ma, and therefore record the denudational response to rifting. An AHe age does not, however, necessarily refer to a distinct geologic event because a range of t - T paths can yield the same AHe age (Wolf et al., 1998). Samples here are likely to have been long exposed to temperatures corresponding to the ^4He PRZ. In sum, because the sampled basement rocks were situated at <2 – 3 km depths and hence shallower than the AFT closure isotherm, cooling histories were complex. More robust data on cooling histories were searched by combined thermal modeling of AFT and AHe data using the HeFTy

program (Ketcham, 2005; see footnote 1) and its multikinetic apatite annealing model. Because AFT annealing kinetics are sensitive to parameter choices, we measured the kinetic parameter D_{par} for each sample (Donelick et al., 1999). D_{par} (in μm) is the mean fission-track etch pit diameter parallel to the c -axis and serves as a proxy for determining chemistry-dependent AFT annealing behavior at crystal level (see footnote 1). Four key requirements guided the forward search for best-fitting t - T paths. These were present-day mean surface temperature ($\sim 30^\circ\text{C}$); an onset of rifting at 25 ± 10 Ma; Eocene paleotemperatures compatible with burial under 0.9 km of sedimentary rocks; and comparatively shallow depths during the Late Cretaceous, because Coniacian layers (91–87 Ma) are capped by bauxite (Roger et al., 1989), which is a surface weathering product. Pre-Cretaceous histories were unknown and therefore left as free parameters.

Despite weak evidence for minor sources of variation due to compositional effects on response to annealing of a few individual apatite grains (Fig. DR1), all the individually modeled samples exhibit relatively coherent thermal histories (Fig. 3), and hence a common geomorphic past with an onset of rapid cooling at 32–20 Ma. Denudation rates (km m.y.^{-1}) are routinely estimated from cooling rates ($^\circ\text{C m.y.}^{-1}$) for a given geothermal gradient ($^\circ\text{C km}^{-1}$). Here, the optimal agreement between the stratigraphy-derived (e.g., 1.75 km; cf. Fig. 2) and LTT-derived denudation depths requires a geothermal gradient of $\geq 33^\circ\text{C km}^{-1}$. Such a value for a craton is high, but nevertheless consistent with a cooler cratonic gradient having locally steepened prior to rifting due to burial beneath the low-conductivity carbonate package. Blanketing increased the geotherm in the sampled underlying rock units while retaining normal background characteristics deeper in the crust.

DISCUSSION

Thermal modeling strategies form a continuum between inverse and forward procedures. With inversion schemes, the search for optimal solutions is loosely guided by a broad t - T parameter space, in which large numbers of forward t - T models are generated randomly and ranked by some statistical goodness-of-fit criterion. However, goodness of fit does not guarantee the geologic uniqueness (and hence accuracy) of solutions, because parameter choices (including the fitness test) determine the outcome as much as the data (Ketcham, 2005). Here, attempts at inverse modeling yielded far less robust results, capturing only the late Cenozoic rapid cooling event with poor precision on timing. Forward modeling is not exempt from nonuniqueness, but in this study the double unknowns of denudation rate and geothermal gradient have been reduced to just the latter because of the good independent information

from the stratigraphy. This could be seen to imply that the success of LTT as a method in quantitative geomorphology is driven by the quality of external inputs and that LTT is therefore prey to circular reasoning. However, the internal consistency of this study also indicates that if most required parameters are available, LTT will predict mean denudation values with some accuracy. The science of geology strives to reconstruct continuous histories with incomplete archives, and so any source of data likely to narrow the range of uncertainty is extremely valuable.

The capacity of LTT to robustly support or challenge geomorphic models of scarp evolution is affected by its limited ability to detect spatial differences in relative denudation depths ≤ 1 km. Here this is apparent in the inability of the data to resolve the coast-to-scarp erosion gradients dictated by the stratigraphy (Fig. 2). Fitting calculated denudation depths to the stratigraphically determined values of Figure 2 would only be possible if different geothermal gradients were assumed for each sample. This seems implausible across such a small geographic area, and would require unrealistically high values (>50 °C km⁻¹) for the less eroded coastal area. Currently, geomorphic models envisage initial scarp geometries either as crustal downwarps, shoulder upwarps, or degrading plateau edges (e.g., Braun and van der Beek, 2004). Downwarps are difficult to detect either in rock packages or in the topography, but at volcanic margins, seaward flexural deformations have been interpreted as synrift rollover structures (Sheth, 1998; Lenoir et al., 2003). In theory, they can also be generated flexurally (Gunnell and Fleitout, 1998) or supported dynamically (Morley and Westaway, 2006) by the effects of post-rift sediment-related and/or cooling-related loads. By applying suitable EET values, the Oman example shows that seaward downwarping can occur and affect a footwall upland (Fig. 2), provided that post-rift erosion and sedimentation are distributed so as to cause maximum flexural uplift in the inner coastal belt.

Unlike Ollier and Pain (1997), who advocated topographic warping of the pre-rift plateau surface and thus its preservation at the coast, we suggest that seaward downwarping is not incompatible with rift-shoulder uplift (e.g., caused by mechanical unloading of the lithosphere during initial tectonic extension; Petit et al., 2007). However, we conceive it as a post-rift phenomenon in which (1) sediment loading occurs offshore, and (2) denudation minima occur inboard of the escarpment (here due to aridity) on a regional scale, and locally near the coast, where the buttes have been preserved as islands of low erosion between catchments expanding into the interior. The resulting geometry mimics topographic warping, but unlike the model proposed by Ollier and Pain, denudation is emphatically not negligible at the coastline.

CONCLUSION

The occurrence of seaward downwarping at rifted margins has been widely discounted on the basis of denudation signatures derived from LTT inversion schemes. However, absence of evidence is not evidence of absence, because AHe and AFT systems often lack the appropriate sensitivity to accurately delimit either post-rift denudation histories or crustal deformation geometries (Braun and van der Beek, 2004; this study). Furthermore, buttes that would testify to plateau-edge cambering may be rare features with low preservation potential, because in non-arid environments they are eroded fully within 30–40 m.y. after rifting. This study emphasizes that LTT is a useful and potentially accurate tool for quantifying long-term denudation rates, but that its power to discriminate between theoretical denudation patterns is substantially strengthened by incorporating stratigraphic data.

ACKNOWLEDGMENTS

We thank Peter van der Beek, Paul Bishop, and an anonymous reviewer for the constructive comments made on this manuscript. The project was partly funded by the Middle-East Basins Evolution (MEBE) program.

REFERENCES CITED

- Bellahsen, N., Fournier, M., d'Acremont, E., Leroy, S., and Daniel, J.-M., 2006, Fault reactivation and rift localization: The northeastern Gulf of Aden margin: *Tectonics*, v. 25, TC1007, doi: 10.1029/2004tc001626.
- Braun, J., and van der Beek, P., 2004, Evolution of passive margin escarpments: What can we learn from low-temperature thermochronology?: *Journal of Geophysical Research*, v. 109, p. F04009, doi: 10.1029/2004JF000147.
- D'Acremont, E., Leroy, S., Beslier, M.-O., Bellahsen, N., Fournier, M., Robin, C., Maia, M., and Gente, P., 2005, Structure and evolution of the eastern Gulf of Aden conjugate margins from seismic reflection data: *Geophysical Journal International*, v. 160, p. 869–890, doi: 10.1111/j.1365-246X.2005.02524.x.
- Donelick, R.A., Ketcham, R.A., and Carlson, W.D., 1999, Variability of apatite fission-track annealing kinetics II: Crystallographic orientation effects: *American Mineralogist*, v. 84, p. 1224–1234.
- Fournier, M., Bellahsen, N., Fabbri, O., and Gunnell, Y., 2004, Oblique rifting and segmentation of the NE Gulf of Aden passive margin: *Geochimistry, Geophysics: Geosystems*, v. 5, p. Q11005, doi: 10.1029/2004GC000731.
- Gallagher, K., Brown, R.W., and Johnson, C.J., 1998, Geological applications of fission track analysis: *Annual Review of Earth and Planetary Sciences*, v. 26, p. 519–572, doi: 10.1146/annurev.earth.26.1.519.
- Gunnell, Y., and Fleitout, L., 1998, Shoulder uplift of the Western Ghats passive margin, India—A flexural model: *Earth Surface Processes and Landforms*, v. 23, p. 391–404, doi: 10.1002/(SICI)1096-9837(199805)23:5<391:AID-ESP853>3.0.CO;2-5.
- Gunnell, Y., Gallagher, K., Carter, A., Widdowson, M., and Hurford, A.J., 2003, Denudation history of the continental margin of western peninsular India during the Mesozoic and Cenozoic: *Earth*

- and Planetary Science Letters, v. 215, p. 187–201, doi: 10.1016/S0012-821X(03)00380-7.
- Ketcham, R.A., 2005, Forward and inverse modeling of low-temperature thermochronometry data: *Reviews in Mineralogy and Geochemistry*, v. 58, p. 275–314, doi: 10.2138/rmg.2005.58.11.
- Lenoir, X., Feraud, G., and Geoffroy, L., 2003, High-rate flexure of the East Greenland volcanic margin: Constraints from ⁴⁰Ar/³⁹Ar dating of basaltic dykes: *Earth and Planetary Science Letters*, v. 673, p. 1–14.
- Leroy, S., Gente, P., Fournier, M., d'Acremont, E., Patriat, P., Beslier, M.-O., Bellahsen, N., Maia, M., Blais, A., Perrot, J., Al-Kathiri, A., Merkouriev, S., Fleury, J.-M., Ruellan, P.-Y., Lepvrier, C., and Huchon, P., 2004, From rifting to spreading in the eastern Gulf of Aden: A geophysical survey of a young oceanic basin from margin to margin: *Terra Nova*, v. 16, p. 185–192, doi: 10.1111/j.1365-3121.2004.00550.x.
- Mercolli, I., Briner, A.P., Frei, R., Schönberg, R., Nägler, T.F., Kramers, J., and Peters, T., 2006, Lithostratigraphy and geochronology of the Neoproterozoic crystalline basement of Salalah, Dhofar, Sultanate of Oman: *Precambrian Research*, v. 145, p. 182–206, doi: 10.1016/j.precamres.2005.12.002.
- Morley, C.K., and Westaway, R., 2006, Subsidence in the super-deep Pattani and Malay basins of Southeast Asia: A coupled model incorporating lower-crustal flow in response to post-rift sediment loading: *Basin Research*, v. 18, p. 51–84, doi: 10.1111/j.1365-2117.2006.00285.x.
- Ollier, C.D., and Pain, C., 1997, Equating the basal unconformity with the palaeoplain: A model for passive margins: *Geomorphology*, v. 19, p. 1–15, doi: 10.1016/S0169-555X(96)00048-7.
- Petit, C., Fournier, M., and Gunnell, Y., 2007, Tectonic and climatic controls on rift escarpments: Erosion and flexural rebound of the Dhofar passive margin (Gulf of Aden, Oman): *Journal of Geophysical Research*, v. 112, B03406, doi: 10.1029/2006JB004554.
- Platel, J.-P., Roger, J., Peters, T.J., Mercolli, I., Kramers, J.D., and Le Métour, J., 1992, Geological map of Salalah, Sultanate of Oman: Oman Ministry of Petroleum and Minerals, Directorate General of Mines, sheet NE 40–09, scale 1:250,000.
- Roger, J., Platel, J.-P., Cavelier, C., and Bourdillon-de-Grisac, C., 1989, Données nouvelles sur la stratigraphie et l'histoire géologique du Dhofar (Sultanat d'Oman): *Bulletin de la Société Géologique de France*, v. 2, p. 265–277.
- Sheth, H.C., 1998, A reappraisal of the coastal Panvel flexure, Deccan traps, as a listric-fault-controlled reverse drag structure: *Tectonophysics*, v. 294, p. 143–149, doi: 10.1016/S0040-1951(98)00148-6.
- Watchorn, F., Nichols, G.J., and Bosence, D.W.J., 1998, Rift-related sedimentation and stratigraphy, southern Yemen (Gulf of Aden), in Purser, B.H., and Bosence, D.W.J., eds., *Sedimentation and tectonics of rift basins: Red Sea–Gulf of Aden*: London, Chapman and Hall, p. 165–189.
- Wolf, R.A., Farley, K.A., and Kass, D.M., 1998, Modeling of the temperature sensitivity of the apatite (U-Th)/He thermochronometer: *Chemical Geology*, v. 148, p. 105–114, doi: 10.1016/S0009-2541(98)00024-2.

Manuscript received 20 December 2006

Revised manuscript received 5 March 2007

Manuscript accepted 11 March 2007

Printed in USA